

Manometric parameters in patients with suspected gastroesophageal reflux disease and normal pH monitoring

Parâmetros manométricos em pacientes com suspeita de doença do refluxo gastroesofágico e pHmetria normal

CAMILLE D. BENATTI¹, FERNANDO A. M. HERBELLA¹, MARCO G. PATTI²

ABSTRACT

Background: pH monitoring is the gold standard test for the evaluation of Gastroesophageal reflux disease (GERD) but esophageal manometry is classically not indicated for making or confirming a suspected diagnosis of GERD. This study aims to evaluate the manometric findings in patients with suspected GERD and normal pH monitoring. **Methods:** 100 adult patients with suspected GERD were retrospectively studied. Patients were divided in Group A: normal reflux score (n=60, 72% women, mean age 51 years) and Group B: abnormal reflux score (n=40, 70% women, mean age 54 years). All patients underwent an upper endoscopy, esophageal manometry and pH monitoring. **Results:** Heartburn was more frequent in group B and epigastric pain was more frequent in the group A, while the prevalence of other symptoms was similar between groups. Abnormal endoscopy, hiatal hernia and esophagitis were more frequent in group B with significant risk for GERD. Lower esophageal sphincter (LES) length and pressure were lower in patients from group B. Esophageal motility was similar between groups. **Conclusions:** Our results show that: (1) symptoms are unreliable to diagnose GERD, (2) abnormal endoscopy is more frequent in patients with GERD, (3) LES length and pressure are decreased in patients with GERD, and (4)

patients with clinical predictors for GERD are not more likely to have manometric parameters to suggest GERD.

Keywords: Gastroesophageal Reflux Disease, Manometry, pH monitoring, Symptoms.

RESUMO

Introdução: A pHmetria é o padrão ouro na avaliação da doença do refluxo gastroesofágico (DRGE), porém, a manometria não é, classicamente, indicada para diagnosticar ou confirmar suspeita de DRGE. Este estudo visa avaliar os achados manométricos em pacientes com DRGE e pHmetria normal. **Método:** Foram estudados retrospectivamente 100 pacientes adultos com suspeita de DRGE. Foram divididos em grupo A: índice de refluxo normal (n=60, 72% mulheres, idade média 51 anos) e grupo B: índice de refluxo anormal (n=40, 70% mulheres, idade média 54 anos). Todos os pacientes fizeram endoscopia, manometria e pHmetria esofágicas. **Resultados:** Pirose foi mais frequente no grupo B e epigastria no grupo A, enquanto que a prevalência dos demais sintomas foi similar entre os grupos. Endoscopia anormal, hérnia de hiato e esofagite foram mais frequentes no grupo B, com risco significante para DRGE. O comprimento e a pressão do esfíncter esofágico inferior foram menores

1. Department of Surgery, Escola Paulista de Medicina, Federal University of Sao Paulo, Sao Paulo, Brazil. 2. Department of Surgery, Pritzker School of Medicine, University of Chicago, Chicago, USA. **Corresponding author:** Fernando A. M. Herbella - Department of Surgery - Escola Paulista de Medicina - Rua Diogo de Faria, 1.087 - cj. 301 - Cep: 04037-003 - São Paulo - SP - Brazil - **e-mail:** herbella.dcir@epm.br. **Received in:** 25/11/2013. **Approved in:** 02/02/2014.

nos pacientes do grupo B. A motilidade esofágica foi similar entre os grupos. **Conclusão:** Nossos resultados mostram que: 1 - os sintomas não são discriminatórios no diagnóstico de DRGE; 2 - endoscopia anormal é mais frequente em pacientes com DRGE; 3 - comprimento e pressão do esfíncter esofágico inferior foram menores em pacientes com DRGE, e 4 - pacientes com preditores clínicos para DRGE não têm necessariamente achados manométricos que possam sugerir a DRGE.

Unitermos: Doença do Refluxo Gastroesofágico, Manometria, pHmetria, Sintomas.

INTRODUCTION

Gastroesophageal reflux disease (GERD) is a very common disease; however, it has a complex physiopathology¹ and a myriad of clinical presentations encompassing esophageal and extra-esophageal symptoms². Thus, the diagnosis of GERD may be difficult in some cases. pH monitoring is the gold standard test for the evaluation of GERD even though a significant percentage of false negative results may be linked to this test³.

Manometry may evaluate changes in the esophageal motility that may be associated to GERD physiopathology, such as the lower esophageal sphincter (LES) competence and esophageal body peristalsis; however, according to the American Gastroenterological Association⁴ and the American College of Gastroenterology⁵, manometry is not indicated for making or confirming a suspected diagnosis of GERD.

This study aims to evaluate the manometric findings in patients with suspected GERD and normal pH monitoring.

METHODS

Population

One-hundred non-selected adult patients that underwent esophageal manometry and ambulatory pH monitoring for suspected GERD were retrospectively studied. Patients were grouped on the basis of the results of ambulatory pH monitoring test, group A: normal reflux score (n=60, 72% women, mean age 51 years) and group B: abnormal reflux score (n=40, 70% women, mean age 54 years).

Patients with previous foregut surgery or primary esophageal motility disorders were excluded from the study. Patients were questioned regarding the presence of symptoms that were divided into the following subgroups: esophageal

symptoms (heartburn, regurgitation); extra-esophageal symptoms (thoracic pain, respiratory symptoms, such as cough and asthma or ear nose and throat symptoms) and gastric symptoms (epigastric pain, gastric bloating or fullness and vomiting)

Work up

All patients were submitted to an upper endoscopy to evaluate the presence of hiatal hernia (HH), esophagitis and Barrett's esophagus.

In 48.3% of group A (29 patients) and 45% of group B (18 patients) a barium esophagram was also performed and the presence of HH and gastroesophageal reflux were assessed.

Manometry

Esophageal manometry was performed in all patients. Medications that interfere with esophageal and gastric motility were discontinued 3 days before the study.

Esophageal manometry was performed with an eight-lumen manometry catheter, continuously perfused by pneumohydraulic capillary infusion system connected to a polygraph at least after 8h of fasting. Position, pressure (at the mean respiratory point) and length of the lower esophageal sphincter (LES) were measured using the station pull-through technique. Esophageal body function was assessed by giving 10 swallows of 5 ml of water at 30-s intervals. Amplitude and propagation of the peristaltic waves were assessed. The data were analyzed by computer, using a dedicated software program.

LES length was considered normal when > 2 cm and abdominal length when > 1 cm. LES pressure was considered normal between 14-36 mmHg. A defective LES was defined as the presence of abnormal LES total length or abnormal abdominal length or hypotonicity. Distal esophageal amplitude (DEA) was considered normal when the mean pressure in distal esophagus (sensor located 3cm above the upper border of the LES) was between 60 and 140 mmHg. DEA was considered hypotensive if < 60 mmHg.

pH monitoring

Esophageal pH monitoring was performed on all patients. Acid suppressing medications were discontinued 14 days before the study. During the study, the patients consumed an unrestricted diet. Ambulatory pH monitoring was performed by placing a pH probe 5cm above the upper border of the manometrically determined LES. The data were incorporated into a composite score (DeMeester score), and a score greater than 14.7 was set as abnormal.

Statistics

The student's t test, Bayesian calculations for diagnostic tests, relative risk, Receiver Operating Characteristic (ROC) and Fisher's test were used when appropriated. A value of *p* was considered significant at the 0.05 level. Variables are expressed as mean ± standard deviation.

Ethics

This study was approved by the Research Ethics Committee of the Sao Paulo Federal University.

RESULTS

Population

Group A and B were comparable according to age and gender (table 1).

Symptomatic Evaluation

Heartburn was more frequent in group B and epigastric pain was more frequent in the group A, while the prevalence of other symptoms was similar between groups. However, only heartburn showed a significant risk for GERD (table 1).

Work up

Abnormal endoscopy, hiatal hernia and esophagitis were more frequent in group B with significant risk for GERD (table 2). Four patients had Barrett's esophagus (all of them from group B).

There was no difference in the presence of abnormal esophagram, hiatal hernia or reflux between groups at the esophagram (table 2).

Manometry

LES length and pressure were lower in patients from group B. Esophageal motility was similar between groups. A defective LES was more frequently found in group B and it was the only manometric parameter to increase the risk for GERD (table 3).

The sensitivity and specificity to diagnostic GERD based on abnormal LES length was 22 and 85%, respectively. The sensitivity and specificity to diagnostic GERD based on a hypotonic LES was 70 and 50%, respectively. The sensitivity and specificity to diagnostic GERD based on a defective LES was 87 and 22%, respectively.

A ROC analysis of the LES pressure as a predictor for GERD showed an area under the curve of 0.6 (95% confidence interval 0.5 to 0.7 *p*=0.02). The sensitivity and specificity for different cutoff values are depicted on figure 1.

Table 1. Demographics and symptoms for groups with and without gastroesophageal reflux disease.

	Group A (GERD -) (n=60)	Group B (GERD +) (n=40)	<i>p</i>	Relative risk (95% confidence interval)
Age (years)	51.5 ± 12.9	53.6 ± 13.73	0.43	NA
Gender (% females)	72	70	0.17	0.9 (0.6-1.6)
Heartburn	33 (55%)	32 (80%)	< 0.01	2.1 (1.1 - 4.1)
Regurgitation	33 (55%)	25 (62.5%)	0.12	1.2 (0.7-2.0)
Epigastric pain	32 (53%)	14 (35%)	0.03	0.6 (0.4-1.1)
Dysphagia	7 (11.6%)	5 (12.5%)	0.24	1.0 (0.5-2.1)
Cough	9 (15%)	8 (20%)	0.17	1.2 (0.7-2.2)
Asthma	2 (3.3%)	2 (5%)	0.17	1.3 (0.5-3.5)
Vomiting	9 (15%)	8 (20%)	0.17	1.2 (0.7-2.2)
Retrosternal pain	10 (16.6%)	7 (17.5%)	0.21	1.0 (0.5-1.9)
Esophageal symptoms	45 (75%)	34 (85%)	0.3	1.5 (0.7-3.1)
Extra-esophageal symptoms	26 (43%)	16 (40%)	0.8	0.9 (0.5-1.5)
Gastric symptoms	54 (90%)	36 (90%)	1	1 (0.4-2.2)

NA: not applicable

Subgroups analysis

The analysis of manometric parameters of subgroups with normal pH monitoring and predictors for gastroesophageal reflux disease showed that the LES pressure is lower in patients with heartburn and an abnormal endoscopy compared to other patients from group A (table 4).

Table 2. Preoperative work up for groups with and without gastroesophageal reflux disease.

	Group A (GERD -) (n=60)	Group B (GERD +) (n=40)	<i>p</i>	Relative risk (95% confidence interval)
Abnormal endoscopy	31 (51.6%)	35 (87.5%)	0.02	3.6 (1.5-8.3)
- Hiatal hernia	20 (33.3%)	28 (70%)	< 0.01	2.5 (1.5-4.4)
- Esophagitis	20 (33.3%)	25 (62.5%)	< 0.01	2.0 (1.2-3.4)
Abnormal esophagram	21 (72.4%)	13 (72.2%)	0.24	0.9 (0.6-1.6)
- Hiatal hernia	21 (72.4%)	11 (61.1%)	0.18	0.8 (0.5-1.4)
- Reflux	8 (27.5%)	7 (38.8%)	0.18	1.2 (0.6-2.2)

NA: not applicable

Table 3. Esophageal manometry for groups with and without gastroesophageal reflux disease.


	Group A (GERD -) (n=60)	Group B (GERD +) (n=40)	p	Relative risk (95% confidence interval)
LES length (cm)	2.2 ± 0.516	2.2 ± 1.055	< 0.01	1.3 (0.8-2.3)
% abnormal	9 (15%)	9 (22.5%)	0.4	
LES abdominal length (cm)	0.7 ± 0.7	0.5 ± 0.6	0.16	1.5 (0.9-2.6)
% abnormal	33 (55%)	28 (70%)	0.10	
LES pressure (mmHg)	15.6 ± 6.6	12.4 ± 4.9	0.05	1.7 (1.0-2.9)
% hypotonic	30 (50%)	28 (70%)	0.06	
Defective LES	39 (65%)	35 (87.5%)	0.02	2.4 (1.1-5.6)
Relaxation pressure of LES (mmHg)	-1.1 ± 1.5	-1.3 ± 1.2	0.35	
Mean wave amplitude (mmHg)	124.0 ± 55.9	102.5 ± 58.4	0.06	1.6 (1-2.6)
hypocontractility	8 (13%)	11 (27.5%)	0.11	
Peristalsis (%)	94.9 ± 12.6	95.7 ± 12.5	0.74	

LES: lower esophageal sphincter

Table 4. Analysis of manometric parameters of subgroups with normal pH monitoring and predictors for Gastroesophageal reflux disease.

	Heartburn + GERD - (n=33)	Abnormal endoscopy GERD - (n=30)	Hiatal hernia (n=20)	Esophagitis (n=20)	Abnormal EDA and Heartburn + (n=17)	Group A (GERD -) (n=60)	Group B (GERD +) (n=40)
Abdominal LES length	6 (18%) p = 0.77 vs GERD - p = 1 vs GERD +	8 (27%) p = 0.25 vs GERD - p = 0.78 vs GERD +	6 (30%) p = 0.18 vs GERD - p = 0.54 vs GERD +	6 (30%) p = 0.18 vs GERD - p = 0.54 vs GERD +	5 (29%) p = 0.28 vs GERD - p = 0.73 vs GERD +	9 (15%)	9 (22%)
Abdominal length	22 (67%) p = 0.37 vs GERD - p = 0.8 vs GERD +	19 (63%) p = 0.5 vs GERD - p = 0.61 vs GERD +	13 (65%) p = 0.6 vs GERD - p = 0.77 vs GERD +	14 (70%) p = 0.3 vs GERD - p = 1 vs GERD +	13 (76%) p = 0.16 vs GERD - p = 0.75 vs GERD +	33 (55%)	28 (70%)
LES pressure	22 (67%) p = 0.13 vs GERD - p = 0.8 vs GERD +	18 (60%) p = 0.5 vs GERD - p = 0.44 vs GERD +	13 (65%) p = 0.3 vs GERD - p = 0.77 vs GERD +	13 (65%) p = 0.3 vs GERD - p = 0.77 vs GERD +	14 (82%) p = 0.02 vs GERD - p = 0.51 vs GERD +	30 (50%)	28 (70%)
Defective LES	26 (79%) p = 0.23 vs GERD - p = 0.35 vs GERD +	21 (70%) p = 0.81 vs GERD - p = 0.12 vs GERD +	14 (70%) p = 0.78 vs GERD - p = 0.15 vs GERD +	15 (75%) p = 0.58 vs GERD - p = 0.27 vs GERD +	14 (82%) p = 0.23 vs GERD - p = 0.68 vs GERD +	39 (65%)	35 (87%)
Hypocontractility	2 (6%) p = 0.48 vs GERD - p = 0.02 vs GERD +	3 (10%) p = 0.74 vs GERD - p = 0.07 vs GERD +	1 (5%) p = 0.43 vs GERD - p = 0.04 vs GERD +	2 (10%) p = 1 vs GERD - p = 0.18 vs GERD +	1 (6%) p = 0.67 vs GERD - p = 0.08 vs GERD +	8 (13%)	11 (27%)

LES=lower esophageal sphincter/GERD = Gastroesophageal reflux disease

Figure 1. Sensitivity and specificity for different cutoff values in a Receiver Operating Characteristic (ROC) curve of the lower esophageal sphincter pressure as a predictor for positive pH monitoring.

DISCUSSION

Our results show that: (1) symptoms are unreliable to diagnose GERD, (2) abnormal endoscopy is more frequently found in patients with GERD, (3) LES length and pressure are decreased in patients with GERD, and (4) patients with clinical predictors for GERD are not more likely to have manometric parameters to suggest GERD.

GERD diagnosis

Different previous publications showed that symptoms are unreliable for the diagnosis of GERD^{6,7}, although the labeling of patients as refluxers based on symptoms questionnaires is still a common practice, even in important scientific publications^{8,9}, in spite of that most of these questionnaires were not validated in comparison to esophageal ambulatory pH monitoring¹⁰. In our series, heart-burn was the only symptom with a mathematical association to GERD; however, more than half of the patients without GERD complaint of heartburn. A thoughtfully investigation of patients with suspect GERD especially with pH monitoring may avoid much improper and costly medical therapy¹¹. Upper digestive endoscopy is quoted as highly specific but with a low sensitivity for the diagnosis of GERD¹². It is well-known that the absence of esophagitis does not preclude the diagnosis of GERD; however, the presence of esophagitis is considered an appanage of refluxers by most authors¹³. In our series, esophagitis was found in one third of the GERD - patients by pH monitoring, a rate ranging from 20 to 47% in other previous studies^{6,14}.

We cannot hypothesize the basis for esophagitis in patients without GERD. It may probably reflect a false negative pH monitoring or a decreased mucosal resistance even to physiologic reflux. Whether these patients (with esophagitis and a negative pH monitoring) should be considered GERD + will depend on a case by case clinical judgment. Similarly, HH was more frequently found in GERD + patients although a significant number of GERD - also presented with this finding. Despite the fact that one third of GERD - patients may have clinical significant findings at the endoscopy, this test showed a high diagnostic yield for GERD and it should be routinely performed in patients with suspected GERD. Barium swallow (esophagram) did not show a good diagnostic test to discriminate GERD.

Ambulatory prolonged pH monitoring is still the gold standard test for the diagnosis of GERD^{13,15} although some argue that impedance-pH should take this position at the podium¹⁶. A false negative ranging from 4 to 7% may be associated to the test in non-selected populations, such as non-erosive GERD or chest pain¹⁷⁻¹⁸ that may be explained by hypersalivation linked to the presence of the transnasal catheter, changes in life style during the period of the test, and daily variability¹⁹. Also, visceral hypersensitivity may lead to GERD in patients with normal esophageal acid exposure²⁰. We intended to demonstrate in the current study whether patients with clinical predictors for GERD would present with esophageal manometry parameters that could identify a false negative pH monitoring.

Manometry and GERD

GERD has a complex physiopathology¹ and the observation of only esophageal manometry parameters may not diagnose the disease. A defective LES is not a guarantee of GERD since other natural antireflux mechanisms may be acting efficiently^{1,21}. On the other side, a manometric normal LES may be found in refluxers since abnormal transient relaxation of the LES may be implicated in the genesis of GERD and it is not easily detected by routine manometry²¹. In our results, although mean LES length and pressure are decreased in patients with GERD and a defective LES leads to a 2.5 increased chance to GERD still a normal LES was found in 13% of the GERD + patients and a defective LES was found in 65% of the GERD - patients.

The prevalence of a defective LES do not suggest pathologic reflux even when a subgroup of GERD - patients with a high likelihood to be real refluxers (abnormal endoscopy and heartburn) are evaluated. Very interestingly too, a ROC curve analysis of the cutoff value for LES pressure showed that the best specificity and sensitivity for GERD diagnosis was found exactly at the current adopted lower limit for LES normal pressure (14 mmHg). The analysis of a different series failed to show differences of the LES between GERD + and GERD - patients⁶. Esophageal peristalsis is also an important component of the antireflux mechanism because it is the main determinant of esophageal clearance of the refluxate⁽¹⁾. We did not find differences in esophageal amplitude or peristalsis in patients GERD - or GERD +, as observed by others as well⁶.

Although esophageal manometry parameters do not diagnose GERD they may grade the disease. A direct correlation between acid exposure and progressive deteriorate degrees of LES and esophageal peristalsis function was demonstrated²². Also, weakened peristalsis and defectives LES has been long associated to more severe GERD and Barrett's esophagus²³⁻²⁵. Esophageal manometry parameters have also been tested as predictors for GERD therapy outcomes, but different studies failed to link motility and outcomes for surgical treatment of GERD²⁵⁻²⁸.

CONCLUSIONS

The current study has some limitations. First, it is a retrospective study with the bias associated to this study design. Second, we only measured acid reflux although we believe that non-acid reflux parallels acid reflux and our results would be unchanged²⁹. Lastly, we did not follow the patients up or evaluate response to antacids since some of them were not treated in the same institution where the esophageal function

tests were performed. Despite these limitations, we showed that esophageal manometry must not be used as a unique tool to diagnosis GERD; however, the finding of a defective LES must be summed to other clinical parameters to allow a correct judgment of GERD diagnosis in difficult cases. Furthermore, esophageal manometry is also a useful tool to correct placement of pH monitoring catheters and allows the diagnosis of unsuspected primary motility disorders. As such, esophageal manometry should be liberally done in patients under evaluation for suspected GERD.

REFERENCES

- Herbella FA, Patti MG. Gastroesophageal reflux disease: From pathophysiology to treatment. *World J Gastroenterol*. 2010 Aug 14;16(30):3745-9.
- Moore JM, Vaezi MF. Extraesophageal manifestations of gastroesophageal reflux disease: real or imagined? *Curr Opin Gastroenterol*. 2010 Jul;26(4):389-94.
- Sweis R, Fox M, Anggiansah A, Wong T. Prolonged, wireless pH-studies have a high diagnostic yield in patients with reflux symptoms and negative 24-h catheter-based pH-studies. *Neurogastroenterol Motil*. 2011 May;23(5):419-26.
- Pandolfino JE, Kahrilas PJ; American Gastroenterological Association. American Gastroenterological Association medical position statement: Clinical use of esophageal manometry. *Gastroenterology*. 2005 Jan;128(1):207-8.
- Hirano I, Richter JE; Practice Parameters Committee of the American College of Gastroenterology. ACG practice guidelines: esophageal reflux testing. *Am J Gastroenterol*. 2007 Mar;102(3):668-85.
- Bello B, Zoccali M, Gullo R, Allaix ME, Herbella FA, Gasparitis A, Patti MG. Gastroesophageal reflux disease and antireflux surgery—what is the proper preoperative work-up? *J Gastrointest Surg*. 2013 Jan;17(1):14-20.
- Chan K, Liu G, Miller L, Ma C, Xu W, Schlachta CM, Darling G. Lack of correlation between a self-administered subjective GERD questionnaire and pathologic GERD diagnosed by esophageal pH monitoring. *J Gastrointest Surg* 2010;14:427-436.
- Liang B, Wang M, Yi Q, Feng Y. Association of gastroesophageal reflux disease risk with exacerbations of chronic obstructive pulmonary disease. *Dis Esophagus*. 2013 Jan 10. doi: 10.1111/dote.12014. [Epub ahead of print]
- Bruley des Varannes S, Ducrotté P, Vallot T, Garofano A, Bardoulat I, Carrois F, Ricci L. Gastroesophageal reflux disease: Impact on work productivity and daily-life activities of daytime workers. A French cross-sectional study. *Dig Liver Dis*. 2013 Mar;45(3):200-6.
- Vakil NB, Halling K, Becher A, Rydén A. Systematic review of patient-reported outcome instruments for gastroesophageal reflux disease symptoms. *Eur J Gastroenterol Hepatol*. 2013 Jan;25(1):2-14.
- Patti MG, Diener U, Tamburini A, Molena D, Way LW. Role of esophageal function tests in diagnosis of gastroesophageal reflux disease. *Dig Dis Sci*. 2001 Mar;46(3):597-602.
- Hatlebakk JG. Endoscopy in gastro-oesophageal reflux disease. *Best Pract Res Clin Gastroenterol*. 2010 Dec;24(6):775-86. doi: 10.1016/j.bpg.2010.09.005.
- Moayyedi P, Talley NJ. Gastro-oesophageal reflux disease. *Lancet*. 2006 Jun 24;367(9528):2086-100.
- Frootan M, Choobtashani S, Azargashb E, Amin PM, Hamide M, Zali MR, Haleh A. Non-erosive reflux disease compared with erosive esophagitis with regards to acid reflux and symptom patterns. *Turk J Gastroenterol*. 2011 Oct;22(5):464-71.
- Herbella FA, Nipominick I, Patti MG. From sponges to capsules. The history of esophageal pH monitoring. *Dis Esophagus*. 2009;22(2):99-103.
- Castell DO. Reflux testing in the 21st century: is there a role for pH only? *Clin Gastroenterol Hepatol*. 2008 Aug;6(8):840-1. doi: 10.1016/j.cgh.2008.04.007. Epub 2008 Jun 27.
- Madan K, Ahuja V, Gupta SD, Bal C, Kapoor A, Sharma MP. Impact of 24-h esophageal pH monitoring on the diagnosis of gastroesophageal reflux disease: defining the gold standard. *J Gastroenterol Hepatol*. 2005 Jan;20(1):30-7.
- Fuchs KH, DeMeester TR, Albertucci M. Specificity and sensitivity of objective diagnosis of gastroesophageal reflux disease. *Surgery*. 1987 Oct;102(4):575-80.
- Ayazi S, Hagen JA, Zehetner J, Banki F, Augustin F, Ayazi A, DeMeester SR, Oh DS, Sohn HJ, Lipham JC, DeMeester TR. Day-to-day discrepancy in Bravo pH monitoring is related to the degree of deterioration of the lower esophageal sphincter and severity of re-flux disease. *Surg Endosc*. 2011 Jul;25(7):2219-23. doi: 10.1007/s00464-010-1529-5.
- Remes-Troche JM. The hypersensitive esophagus: pathophysiology, evaluation, and treatment options. *Curr Gastroenterol Rep*. 2010 Oct;12(5):417-26. doi: 10.1007/s11894-010-0122-3.
- Hershovici T, Mashimo H, Fass R. The lower esophageal sphincter. *Neurogastroenterol Motil*. 2011 Sep;23(9):819-30. doi: 10.1111/j.1365-2982.2011.01738.x. Epub 2011 Jun 29.
- Lee J, Anggiansah A, Anggiansah R, Young A, Wong T, Fox M. Effects of age on the gastroesophageal junction, esophageal motility, and reflux disease. *Clin Gastroenterol Hepatol*. 2007 Dec;5(12):1392-8.
- Savarino E, Gemignani L, Pohl D, Zentilin P, Dulbecco P, Assandri L, Marabotto E, Bon-fanti D, Inferrera S, Fazio V, Malessi A, Tutuian R, Savarino V. Oesophageal motility and bolus transit abnormalities increase in parallel with the severity of gastro-oesophageal re-flux disease. *Aliment Pharmacol Ther*. 2011 Aug;34(4):476-86. doi: 10.1111/j.1365-2036.2011.04742.x. Epub 2011 Jun 14.
- Ang D, Blondeau K, Sifrim D, Tack J. The spectrum of motor function abnormalities in gastroesophageal reflux disease and Barrett's esophagus. *Digestion*. 2009;79(3):158-68. doi: 10.1159/000210265.
- Lord RV, DeMeester SR, Peters JH, Hagen JA, Elyssnia D, Sheth CT, DeMeester TR. Hiatal hernia, lower esophageal sphincter incompetence, and effectiveness of Nissen fundoplication in the spectrum of gastroesophageal reflux disease. *J Gastrointest Surg*. 2009 Apr;13(4):602-10. doi: 10.1007/s11605-008-0754-x.
- Patti MG, Perretta S, Fisichella PM, D'Avanzo A, Galvani C, Gorodner V, Way LW. Laparoscopic antireflux surgery: preoperative lower esophageal sphincter pressure does not affect outcome. *Surg Endosc*. 2003 Mar;17(3):386-9.
- Patti MG, Robinson T, Galvani C, Gorodner MV, Fisichella PM, Way LW. Total fundoplication is superior to partial fundoplication even when esophageal peristalsis is weak. *J Am Coll Surg*. 2004 Jun;198(6):863-9; discussion 869-70.
- Riedl O, Gadenstätter M, Lechner W, Schwab G, Marker M, Ciofica R. Preoperative lower esophageal sphincter manometry data neither impact manifestations of GERD nor outcome after laparoscopic Nissen fundoplication. *J Gastrointest Surg*. 2009 Jul;13(7):1189-97. doi: 10.1007/s11605-009-0890-y. Epub 2009 Apr 16.
- Herbella FA. Critical analysis of esophageal multichannel intraluminal impedance monitoring 20 years later. *ISRN Gastroenterol*. 2012;2012:903240